



# Tidsskriftet

DEN NORSKE LEGEFORENING

## Acute poisoning from substance abuse of benzodiazepine

---

### SHORT REPORT

JON ANDREAS ANDERSSON

Faculty of Medicine University of Oslo  
and  
Valnesfjord Medical Centre  
Fauske Municipality  
Jon Andreas Andersson is a general practitioner.  
The author has completed the ICMJE form and declares no conflicts of interest.

METTE BREKKE

General Practice Research Unit  
University of Oslo  
Professor Mette Brekke is a specialist in general practice.  
The author has completed the ICMJE form and declares no conflicts of interest.

ODD MARTIN VALLERSNES

E-mail: o.m.vallersnes@medisin.uio.no  
Department of General Practice  
University of Oslo  
and  
Department of Emergency General Practice  
Oslo Accident and Emergency Outpatient Clinic  
City of Oslo Health Agency  
Odd Martin Vallersnes is associate professor and a consultant general practitioner.  
The author has completed the ICMJE form and declares no conflicts of interest.

---

### BACKGROUND

Benzodiazepines are also used as intoxicants. This can be dangerous, particularly in multi-substance abuse. We describe cases of acute poisoning from substance abuse of benzodiazepines in patients at the main A&E clinic in Oslo.

### MATERIAL AND METHOD

We included all patients treated for substance abuse poisoning with benzodiazepines and/or z-hypnotics at the Oslo Accident and Emergency Clinic from 1 October 2013 to 30 September 2015. The patients were found through a retrospective review of the A&E clinic's registers. Data from patient records. Diagnosis of the toxic agent was based on the attending doctor's recorded clinical evaluation.

### RESULTS

Of 1 037 cases, 787 (76 %) were men. The median age was 36 (interquartile range 28–46, range 14–78). Clonazepam (Rivotril) was the most frequent drug, with 575 cases (55 %), followed by diazepam (Stesolid, Valium, Vival) 158 (15 %), alprazolam (Xanor) 125 (12 %) and oxazepam (Sobril) 90 (8 %). Imovane, Zopitin and zolpidem (Stilnoct) occurred rarely, in 25 (2 %) and 11 (1 %) cases, respectively. Benzodiazepines were combined with other substances in 936 (90 %) cases, most frequently heroin 484 (47 %), ethanol 321 (31 %) and amphetamine 199 (19 %).

### INTERPRETATION

In substance abuse poisoning, benzodiazepines were very often combined with other intoxicants, most frequently opioids, ethanol and/or

---

### MAIN FINDINGS

Clonazepam (Rivotril), diazepam (Stesolid, Valium, Vival) and alprazolam (Xanor) were the most commonly occurring benzodiazepines in acute poisoning.

In nine out of ten cases, benzodiazepines had been taken in combination with other intoxicants.

---

The use of benzodiazepines has increased sharply since they were first synthesised in the 1950s, and they have gradually replaced barbiturate, hypnotic and anti-anxiety drugs (1). In 2008, 6 % of the Norwegian population was prescribed at least one benzodiazepine (2). Since the 1990s, drugs such as zopiclone (Imovane, Zopitin) and zolpidem (Stilnoct), have taken over for insomnia (3).

Benzodiazepines and z-hypnotics are chemically dissimilar, but both act by increasing the effect of the neurotransmitter gamma-aminobutyric acid on GABA<sub>A</sub> receptors in the brain. This reduces neuronal excitability, with a hypnotic, anxiolytic and anticonvulsive effect as a result (4,5).

Benzodiazepines can also be used as intoxicants, and have adverse long-term effects in the form of impaired cognitive abilities and risk of overdose. Overdosing results in altered level of consciousness, impaired coordination, amnesia and respiratory depression. The risk of an overdose is increased in combination with other intoxicants with depressant effects. Flumazenil is an antidote, but has a short half-life and may cause seizures (4,5).

Z-hypnotics cause less intoxication and entail less risk of inducing dependence (5), but the mechanism is otherwise so similar to benzodiazepines that we have chosen to include them in this study. In the following we use benzodiazepines as a collective term.

12 % of benzodiazepine users account for 59 % of prescribed use in Norway (6). A Swiss study found that heavy users often alternate between strategies to acquire benzodiazepines (7). In an anonymous British questionnaire survey (8), sources of benzodiazepines were general practitioners and family (40 %), online purchasing (27 %), street sales (20 %) and abroad (11 %). Multiple sources were used by 31 %.

We describe acute poisonings treated by the main A&E clinic in Oslo, due to substance abuse of benzodiazepines, with emphasis on the type of benzodiazepine and other intoxicants they were taken in combination with.

## Material and method

We included all cases of substance abuse poisoning with benzodiazepines treated by the main A&E clinic in Oslo (the Oslo Accident and Emergency Outpatient Clinic (OAEOC)), and retrospectively reviewed cases registered by the reception nurse in the patient registration lists in the A&E electronic record system in the period October 2013–September 2015. Poisonings with suicidal intention were not included.

From patient records we noted age, sex, toxic agent, whether the patient was brought by ambulance, observation period, treatment, disposition of symptoms and signs, in accordance with a data registration template developed by the research network European Drug Emergency (DEN) (9). Diagnosis of the toxic agent was based on the attending doctor's assessment as noted in the records, which in turn was based on the patient and/or supplied by third parties, and on clinical findings. No toxicological laboratory diagnostic work was performed.

The project was carried out as a quality improvement study and was approved by the Information Security and Privacy Office at Oslo University Hospital. SPSS version 25 was used to perform analyses. We used the chi-squared test to compare categorical variables.

## Results

Benzodiazepines were involved in 1 037 substance abuse poisonings in the course of a two-year period. The median age of patients was 36 years, range 28–46, range 14–78, and 787 (76 %) were men.

Clonazepam (Rivotril) occurred most frequently, with 575 (55 %) cases, followed by diazepam (Stesolid, Valium, Vival) 158 (15 %) and alprazolam (Xanax) 104 (10 %) (Table 1).

**Table 1**

Combinations of benzodiazepines (including z-hypnotics) and other intoxicants in acute poisoning cases treated at the main A&E clinic in Oslo, 1.10.2013–30.09.2015. Data are stated as number (%). The total number may be higher than the sums in columns or rows, as several benzodiazepines could be combined with the same intoxicant and several intoxicants could be combined with the same benzodiazepine. GHB = gamma hydroxybutyric acid; MDMA = 3,4-methylenedioxymethamphetamine

	Heroin	Heroin + Methadone	Buprenorphine	Other opioids	Amphetamine	Cocaine	MDMA	GHB	Cannabis	Ethanol	Other / unknown	Only benzo
Clonazepam (Rivotril)	318 (66)	28 (62)	23 (68)	21 (29)	131 (66)	18 (56)	10 (56)	31 (63)	62 (65)	159 (50)	10 (43)	
Diazepam (Stesolid, Valium, Vival)	48 (10)	8 (18)	1 (3)	10 (14)	24 (12)	5 (16)	2 (11)	8 (16)	18 (19)	82 (26)	6 (26)	
Alprazolam (Xanax)	77 (16)	4 (9)	6 (18)	7 (10)	27 (14)	6 (19)	3 (17)	5 (10)	10 (11)	23 (7)	1 (4)	
Oxazepam (Sobril)	28 (6)	4 (9)	3 (9)	8 (11)	14 (7)	2 (6)	2 (11)	3 (6)	8 (8)	35 (11)	3 (13)	
Flunitrazepam (Rohypnol)	14 (3)	1 (2)	1 (3)	2 (3)	7 (4)	1 (3)	2 (11)	4 (8)	4 (4)	7 (2)	-	
Zopiclone (Imovane, Zopitin)	3 (1)	-	-	2 (3)	-	-	-	-	2 (2)	13 (4)	-	
Nitrazepam (Apodorm, Mogadon)	7 (1)	-	1 (3)	2 (3)	2 (1)	1 (3)	-	2 (4)	-	5 (2)	-	
Zolpidem (Stilnoct)	-	-	-	2 (3)	-	-	-	-	-	4 (1)	-	
Flurazepam	-	1 (2)	-	-	-	-	-	-	-	-	-	
Unspecified	62 (13)	2 (4)	2 (6)	26 (36)	22 (11)	3 (9)	1 (6)	7 (14)	7 (7)	35 (11)	5 (22)	
<b>Total</b>	<b>484 (100)</b>	<b>45 (100)</b>	<b>34 (100)</b>	<b>72 (100)</b>	<b>199 (100)</b>	<b>32 (100)</b>	<b>18 (100)</b>	<b>49 (100)</b>	<b>95 (100)</b>	<b>321 (100)</b>	<b>23 (100)</b>	

Benzodiazepines were combined with other intoxicants in 936 cases (90 %), most frequently heroin 484 (47 %), ethanol 321 (31 %) and amphetamine 199 (19 %).

Patients were brought to the A&E clinic by ambulance in 569 cases (55 %). On admission 47 patients (5 %) were comatose (Glasgow Coma Scale 8), 619 (60 %) somnolent (GCS score 9–14) and 111 (11 %) had a respiratory rate < 12/min. During the course, 150 (14 %) patients were agitated, experienced anxiety and 41 (4 %) were hypotensive. Naloxone (antidote against opioids) was administered in 187 cases (18 %), sedation in 17 cases (1.7 %), flumazenil in 2 (0.2 %). The median observation period was 4 h 25 min (interquartile range 2 h 30 min – 6 h 11 min, range 3 min – 36 h 53 min (16 %)) the patient was hospitalised in a somatic hospital, and in 20 (2 %) in a psychiatric unit. No patients died at the A&E clinic.

A respiratory rate of < 12/min was more common when benzodiazepines were combined with opioids, 101/601 (17 %) cases vs 10/436 (2 %) (p < 0.001). Naloxone was administered more frequently for this combination: 171/601 (28 %) cases vs 16/436 (4 %) (p < 0.001).

## Discussion

In substance abuse poisonings, benzodiazepines were very often taken in combination with other intoxicants – probably to enhance the undesirable side effects or relieve abstinence symptoms. This is in line with other European studies based on a clinical assessment of toxic cases. Even more cases prove to involve benzodiazepines, when clinical assessments are supported by laboratory analyses (12).

Clonazepam was the dominant toxic agent. This stands in contrast to the Norwegian prescription pattern. Z-hypnotics accounted in 2014 for 27 % of prescriptions, and 50 % more of both diazepam and oxazepam was prescribed than clonazepam, and almost the same amount of nitrazepam and clonazepam (3). However, police seizures clearly reflect the distribution of the most frequently occurring benzodiazepines in substance abuse poisonings. Clonazepam accounted for 52 % of the number of benzodiazepine seizures and 47 % of the quantity seized in 2014, followed by diazepam (27 %), alprazolam (13 % and 18 %), while the quantity of z-hypnotics seized was minimal (13). Z-hypnotics seldom occur in substance abuse poisonings because the intoxicant effect is relatively limited (5). Although Lyphout et al. found that national prescription patterns correlated with the benzodiazepines in substance abuse poisonings (11), the intoxication potential and availability on the illegal market appear to be of greater importance than prescription by doctors.

Altered level of consciousness was the most prominent clinical finding, as expected (4,5). Patients with assumed substance abuse intoxication were systematically at the OAEOC according to a locally developed procedure (14). Antidote is administered only in cases of respiratory depression with respiratory depression had taken opioids in addition. Naloxone is a safer antidote than flumazenil and is therefore administered first. Flumazenil was only administered in two cases shows that naloxone was virtually always sufficient and that respiratory depression is rare in benzodiazepines alone. If the respiratory rate does not improve, or if the patient has not woken within four hours and can give an account of the poisoning, the patient is sent on to hospital. Although the OAEOC has good observation possibilities and a short distance to hospital, we believe that our observations might be applicable to other A&E clinics with observation possibilities, and to hospital emergency departments.

#### STRENGTHS AND WEAKNESSES

Toxicological screening is not carried out at the OAEOC. Diagnoses of toxic agents were accordingly based mainly on self-reporting, which may be incorrect owing to patients' altered level of consciousness, tablets of unknown content from street and online sales, unwillingness to cooperate, or fear of legal reprisals. There may also be variation in doctors' record-keeping and in the inclusion and registration practice of data collected.

Although the majority of substance abuse poisonings in Oslo are treated at the A&E clinic, the sickest patients are taken directly to hospital (10). The picture of benzodiazepine poisonings in the capital is therefore not complete. Nor does the study cover benzodiazepine poisoning without intent, a far larger share of which are sent on from the A&E Clinic to hospital (10). Our material is five years old, and the prevalence of the combination of benzodiazepines and other intoxicants may have changed since then.

#### CONCLUSION

Benzodiazepines are most often taken in combination with other intoxicants in substance abuse poisonings, most frequently heroin, ethanol, and amphetamine. Clonazepam was the most prevalent benzodiazepine.

---

*The article has been peer-reviewed.*

---

#### LITERATURE

1. Wick JY. The history of benzodiazepines. *Consult Pharm* 2013; 28: 538–48. [PubMed][CrossRef]
2. Handal M, Skurtveit S, Mørland JG. Samtidig bruk av ulike benzodiazepiner. *Tidsskr Nor Legeforen* 2012; 132: 526–30. [PubMed][CrossRef]
3. Folkehelseinstituttet. Reseptregisteret. <http://www.reseptregisteret.no> Accessed 3.12.2019.
4. Gaudreault P, Guay J, Thivierge RL et al. Benzodiazepine poisoning. Clinical and pharmacological considerations and treatment. *Drug Saf* 1999; 6: 247–60. [CrossRef]
5. Gunja N. The clinical and forensic toxicology of Z-drugs. *J Med Toxicol* 2013; 9: 155–62. [PubMed][CrossRef]
6. Rossow I, Bramness JG. The total sale of prescription drugs with an abuse potential predicts the number of excessive users: a national prescription data analysis. *Public Health* 2015; 15: 288. [PubMed][CrossRef]
7. Liebrecht M, Schneider M, Buadze A et al. High-dose benzodiazepine dependence: a qualitative study of patients' perceptions on initiation, reasons for relapse, and treatment. *PLoS One* 2015; 10: e0142057. [PubMed][CrossRef]
8. Kapil V, Green JL, Le Lait C et al. Misuse of benzodiazepines and Z-drugs in the UK. *Br J Psychiatry* 2014; 205: 407–8. [PubMed][CrossRef]
9. Wood DM, Heyerdahl F, Yates CB et al. The European Drug Emergencies Network (Euro-DEN). *Clin Toxicol (Phila)* 2014; 52: 239–41. [PubMed][CrossRef]
10. Vallersnes OM, Jacobsen D, Ekeberg Ø et al. Patients presenting with acute poisoning to an outpatient emergency clinic: a one-year observational study. *BMC Emerg Med* 2015; 15: 18. [PubMed][CrossRef]
11. Lyphout C, Yates C, Margolin ZR et al. Presentations to the emergency department with non-medical use of benzodiazepines and Z-drugs: profiling and risk factors. *Eur J Clin Pharmacol* 2019; 75: 77–85. [PubMed][CrossRef]
12. Vallersnes OM, Persett PS, Øiestad EL et al. Underestimated impact of novel psychoactive substances: laboratory confirmation of recreational drug toxicity in Norway. *Clin Toxicol (Phila)* 2017; 55: 636–44. [PubMed][CrossRef]
13. KRIPOS. Narkotika- og dopingstatistikk 2014. Oslo: Politiet, 2015.
14. Vallersnes OM, Jacobsen D, Ekeberg Ø et al. Outpatient treatment of acute poisoning by substances of abuse: a prospective observational cohort study. *Resusc Emerg Med* 2016; 24: 76. [PubMed][CrossRef]

---

Publisert: 29 June 2020. *Tidsskr Nor Legeforen*. DOI: 10.4045/tidsskr.20.0035

Received 11.1.2020, first revision submitted 31.3.2020, accepted 6.5.2020.

Published under open access CC BY-ND. Downloaded from [tidsskriftet.no](http://tidsskriftet.no) 31 March 2023.