

## Puberty starts earlier

---

### OPINIONS

INGVILD SÆRVOLD BRUSERUD

[ingvild.bruserud@uib.no](mailto:ingvild.bruserud@uib.no)

Ingvild Særvold Bruserud PhD, postdoctoral fellow and registered nurse at the Department of Paediatrics, Haukeland University Hospital, Bergen, Norway.

The author has completed the ICMJE form and declares no conflicts of interest.

NINNIE HELÉN BAKKEN OEHME

Ninnie Helén Bakken Oehme, PhD, specialty registrar in paediatrics at the Department of Paediatrics, Haukeland University Hospital, Bergen, Norway.

The author has completed the ICMJE form and declares no conflicts of interest.

ANDRÉ MADSEN

André Madsen PhD, molecular biologist and postdoctoral fellow at the Hormone Laboratory, Department of Medical Biochemistry and Pharmacology, Haukeland University Hospital, Bergen, Norway.

The author has completed the ICMJE form and declares no conflicts of interest.

KAREN ROSENDAHL

Karen Rosendahl, senior consultant radiologist at the Department of Radiology, University Hospital of North Norway and professor at UiT the Arctic University of Norway, Tromsø, Norway.

The author has completed the ICMJE form and declares no conflicts of interest.

ROBERT BJERKNES

Robert Bjercknes, specialist paediatrician and senior consultant at the Department of Paediatrics, Haukeland University Hospital, and Vice Rector for the University of Bergen, Bergen, Norway.

The author has completed the ICMJE form and declares no conflicts of interest.

#### MATHIEU ROELANTS

Mathieu Roelants, PhD, biostatistician and postdoctoral fellow at the Department of Public Health and Primary Care, University of Leuven, Belgium.

The author has completed the ICMJE form and declares no conflicts of interest.

#### PÉTUR B. JÚLÍUSSON

Professor Pétur B. Júlíusson, specialist paediatrician, senior consultant at the Department of Paediatrics, Haukeland University Hospital, Head of the Department for Health Registry Research and Development, Norwegian Institute of Public Health, and professor at the University of Bergen, Norway.

The author has completed the ICMJE form and declares no conflicts of interest.

---

## **New Norwegian data show that the median age for the onset of puberty in Norwegian girls is six months earlier than previously believed.**

New growth curves for Norwegian girls and boys were published in 2009 [\(1\)](#). At that point, there were no data available for the pubertal development of children and adolescents in Norway. Puberty reference data from Denmark collected in the period 1991–1993 were therefore included in Norwegian growth charts for the 4–19 age group [\(2\)](#). Danish and other international studies have later shown that the onset of puberty is shifting towards an earlier age, particularly in girls [\(3\)](#). Knowledge about the pubertal development in Norwegian children is now available to us, thanks to data from the second Bergen Growth Study [\(4, 5\)](#). We have therefore revised the growth curves from 2009 by adding these reference data for normal puberty in girls and boys.

---

## **Using the puberty reference data**

An assessment of pubertal development is generally based on clinical examination of secondary sexual characteristics. In girls, the onset of puberty is defined by whether breast development has started. In boys, the onset of

puberty is defined by whether the testicular volume has reached 4 mL, and this is assessed by comparative palpation of the testes using a Prader orchidometer (see appendix 1 for the clinical methods used to assess pubertal development).

The puberty reference data are based on the so-called Tanner scale, which was published by the paediatrician James Tanner in the 1960s (6, 7). The Tanner scale is used to define the stages of pubertal maturity, with quantity of pubic hair (in both sexes) and breast maturation (in girls) being measured on a scale from 1 to 5. Not all girls/women reach stage B5/P5, and not all boys reach P5, so these stages have been excluded from the reference data. The revised reference data also present new percentiles for first menstruation (menarche).

In the second Bergen Growth Study, ultrasound was used as an additional means of establishing pubertal development. This is a new method used to characterise the stages of breast maturation and testicular volume. Because ultrasound is not available in normal clinical practice, these measurements are not included in the updated reference data.

As before, reference data for pubic hair, breast development, menarche and testicular volume are shown as percentiles on horizontal lines along the x-axis of the growth curves (see appendices 2 and 3). The 3<sup>rd</sup>, 10<sup>th</sup>, 25<sup>th</sup>, 50<sup>th</sup>, 75<sup>th</sup> and 97<sup>th</sup> age percentiles are indicated with numbers. The 3<sup>rd</sup> percentile indicates the age at which 3 % of girls and boys respectively have reached a certain stage.

---

## Clinical consequences

The median age for the *onset* of puberty in Norwegian girls is currently 10.4 years, which is six months earlier than in the Danish reference data used previously. The onset of puberty in Norwegian boys is currently 11.7 years, which matches the Danish reference data (11.8 years). The median age for menarche is 12.7 years in the Norwegian reference data. This is a difference of 8.7 months compared to the Danish reference data, and it is five months earlier than in the first Bergen Growth Study (2003–2006). We have previously recommended that the threshold age for precocious and delayed puberty should be retained in line with current paediatric guidelines (8, 9). Precocious and delayed pubertal development is defined by the ages that are  $\pm 2$  standard deviations from the mean or median. It is recommended that girls under the age of 8 and boys under the age of 9 who present with signs of pubertal development should be assessed with a view to establishing the underlying cause of precocious puberty and possible treatment. According to the new reference data, girls should have started puberty by 13 years of age and boys by 14.5 years of age.

***«Knowledge about the pubertal development in Norwegian children is now available to us, thanks to data from the second Bergen Growth Study»***

The puberty reference data tell us how girls and boys are currently developing in Norway. The references are based on data from a random sample of healthy children (703 girls and 514 boys) aged between 6 and 16 from seven schools in the municipality of Bergen. The study population's prevalence of overweight and obesity is similar to that of the general population of children in Norway, and the relationship between those of Norwegian and non-Norwegian origin is comparable to that found in children and adolescents in Norway as a whole (16.2 % of the girls and 22.6 % of the boys were of non-Norwegian origin). We are therefore of the opinion that the sample is representative of the general Norwegian population in this age group, and that the reference data are applicable to children and adolescents residing in Norway. The reference data will indicate if a puberty marker has appeared early, normally or late, but they provide no information about health or disorders. A thorough clinical examination of the patient and their medical history should be the first step towards establishing precocious or delayed puberty.

---

## REFERENCES

1. Júlíusson PB, Roelants M, Eide GE et al. Vekstkurver for norske barn. Tidsskr Nor Legeforen 2009; 129: 281–6. [PubMed][CrossRef]
2. Juul A, Teilmann G, Scheike T et al. Pubertal development in Danish children: comparison of recent European and US data. Int J Androl 2006; 29: 247–55, discussion 286 - 90. [PubMed][CrossRef]
3. Aksglaede L, Sørensen K, Petersen JH et al. Recent decline in age at breast development: the Copenhagen Puberty Study. Pediatrics 2009; 123: e932–9. [PubMed][CrossRef]
4. Oehme NHB, Roelants M, Saervold Bruserud I et al. Reference data for testicular volume measured with ultrasound and pubic hair in Norwegian boys are comparable with Northern European populations. Acta Paediatr 2020; 109: 1612–9. [PubMed][CrossRef]
5. Bruserud IS, Roelants M, Oehme NHB et al. References for ultrasound staging of breast maturation, Tanner breast staging, pubic hair and menarche in Norwegian girls. J Clin Endocrinol Metab 2020; 105: 1599–607. [PubMed][CrossRef]
6. Marshall WA, Tanner JM. Variations in pattern of pubertal changes in girls. Arch Dis Child 1969; 44: 291–303. [PubMed][CrossRef]
7. Marshall WA, Tanner JM. Variations in the pattern of pubertal changes in boys. Arch Dis Child 1970; 45: 13–23. [PubMed][CrossRef]
8. Oehme N, Bruserud IS, Madsen A et al. Starter puberteten tidligere enn før? Tidsskr Nor Legeforen 2020; 140. doi: 10.4045/tidsskr.20.0043. [PubMed][CrossRef]
9. Norsk barnelegeforening. Pедиатriveileder. 2.4: Normal pubertet. <https://www.helsebiblioteket.no/pediатriveiledere?>

menuitemkeylev1=5962&menuitemkeylev2=5964&key=144406 Accessed  
6.4.2021.

---

Publisert: 24 October 2022. Tidsskr Nor Legeforen. DOI: 10.4045/tidsskr.21.0210

Received 17.3.2021, accepted 6.4.2021.

Copyright: © Tidsskriftet 2026 Downloaded from tidsskriftet.no 25 June 2026.