



Tidsskriftet

DEN NORSKE LEGEFORENING

New international recommendations for type 2 diabetes – consequences for treatment practice in Norway?

PERSPECTIVES

KÅRE I. BIRKELAND

k.i.birkeland@medisin.uio.no

Kåre I. Birkeland, specialist in internal medicine and endocrinology, professor at the University of Oslo and senior consultant at the Section of Renal Medicine, Department of Transplantation Medicine, Oslo University Hospital, Rikshospitalet. He led the work on the treatment of hyperglycaemia for the Norwegian Directorate of Health's national clinical guidelines for diabetes and chairs the Norwegian Diabetes Association's medical committee.

The author has completed the ICMJE form and declares the following conflicts of interest: He has given talks and chaired meetings for companies that market equipment and medication for the treatment of diabetes, such as Astra Zeneca, Boehringer Ingelheim, Lilly, Merck, Novo Nordisk, Roche and Sanofi, with fees paid to the employer and used for own research. He receives a fee for his role as chair of the Norwegian Diabetes Association's medical committee.

SONDRE MELING

Sondre Meling, specialist in internal medicine and endocrinology, senior consultant at the Department of Endocrinology, Stavanger University Hospital, research fellow at the University of Bergen and chair of the Norwegian Endocrine Society's interest group for diabetes, obesity and metabolism.

The author has completed the ICMJE form and declares the following conflicts of interest: He has received fees for chairing meetings and giving talks for several companies that market equipment and medication for the treatment of diabetes, such as Lilly, Novo Nordisk, Boehringer Ingelheim and Sanofi.

INGVILD VATTEN ALSNES

Ingvild Vatten Alsnes, specialist in family medicine and general practitioner in Sandnes. She is an associate professor at the University of Stavanger, a board member of the Norwegian College of General Practice and a member of the research committee of the Norwegian Medical Association. The author has completed the ICMJE form and declares no conflicts of interest.

EMELIE SVENSSON

Emelie Svensson, specialty registrar in family medicine, general practitioner in Alta, clinical director at Alta sexual assault reception centre, Finnmark Hospital Trust, and a member of the Norwegian Diabetes Association's medical committee.

The author has completed the ICMJE form and declares no conflicts of interest.

LARS GULLESTAD

Lars Gullestad, specialist in internal medicine and cardiology, senior consultant at the Department of Cardiology, Oslo University Hospital, Rikshospitalet, and professor emeritus at the University of Oslo. The author has completed the ICMJE form and declares the following conflict of interest: He has received speaker fees from several companies that market medication for the treatment of diabetes, such as Boehringer Ingelheim, AstraZeneca and MSD.

TROND G. JENSSEN

Trond G. Jenssen, specialist in internal medicine and nephrology, senior consultant at the Section of Renal Medicine, Department of Transplantation Medicine, Oslo University Hospital, Rikshospitalet, and professor at the University of Oslo. He led the work on renal medicine for the Norwegian Directorate of Health's national clinical guidelines for diabetes and is the medical advisor for the Norwegian Diabetes Association.

The author has completed the ICMJE form and declares the following conflicts of interest: He has received fees for giving talks and participating in advisory committees from several companies that market medication for the treatment of diabetes, such as Boehringer Ingelheim, AstraZeneca, Novo Nordisk and Bayer and Abbot, and he receives a fee for his role as medical advisor for the Norwegian Diabetes Association.

New international recommendations for the use of anti-hyperglycaemic drugs in patients with type 2 diabetes are at odds with current practice in Norway and the Norwegian Directorate of Health's national clinical guidelines.



Illustration: Andreas Samuelsson

New knowledge indicates that changes are needed in the traditional treatment of patients with type 2 diabetes. Anti-hyperglycaemic drugs that belong to the SGLT2 inhibitor groups (sodium-glucose cotransporter-2 inhibitors) and GLP-1 receptor agonists (glucagon-like peptide-1 receptor agonists) have been shown to reduce cardiovascular events, progression of kidney disease and mortality (1-3). Weight reduction as a result of intensive caloric restriction or bariatric surgery can lead to remission of diabetes in a significant number of patients (4,5), and is proposed as a new primary treatment goal (6). In addition, new medications will soon be available in Norway which show a documented reduction in blood glucose levels and, in particular, a reduction in body weight that far surpasses previous findings (7,8).

«Weight reduction as a result of intensive caloric restriction or bariatric surgery can lead to remission of diabetes in a significant number of patients, and is proposed as a new primary treatment goal»

Several international clinical guidelines with relevance for doctors and patients in Norway have now been revised based on this knowledge (9–12). The European Association for the Study of Diabetes (EASD) and the American Diabetes Association (ADA) recently published updated consensus recommendations for the management of type 2 diabetes (9). The aim of the treatment is to prevent complications and optimise quality of life. The recommendations emphasise a holistic, person-centred approach to managing type 2 diabetes and the importance of good communication between healthcare professionals and patients and of using language that is non-stigmatising and based on facts (e.g. not using words such as *diabetic* and *non-compliant*). Emphasis is placed on combating health inequalities and promoting universal access to treatment. The recommendations also have a strong focus on the importance of weight reduction as a cornerstone of treatment, and a summary is given of new knowledge generated from several cardiorenal outcome studies.

The most important changes in the pharmacological treatment of hyperglycaemia include the recommendation that SGLT2 inhibitors or GLP1 receptor agonists 'should' be prescribed for people with established cardiovascular disease and 'could' be prescribed for patients at high risk of such disease, *regardless* of their HbA1c level and of whether they are already using metformin. About one-third of patients with type 2 diabetes have a diagnosed cardiovascular disease. However, the number in the risk group depends on the criteria applied.

For patients with obesity, a weight reduction of 10–15% is recommended as an important part of the treatment (the current recommendation is 5–10%). This can normalise blood glucose levels without the need for medication in a significant proportion of patients, but will mean that many patients will require more intensive intervention than they currently receive. The DiRECT study showed that 46% of patients with established type 2 diabetes were normoglycaemic without medication one year after participating in an intensive low-calorie weight management programme within primary care (4). Several studies, including a randomised controlled trial conducted in Norway (5), have shown remission in 60–70% of patients after bariatric surgery for morbid obesity. Recently published studies using semaglutide 2.4 mg/week (7) and tirzepatide (8), which are expected to be available in Norway soon, show an average weight reduction of around 10%, and a large proportion of patients achieved a weight reduction in excess of 15%. We believe that a broad and open debate is now needed on the use of and funding for such medication, involving users, GPs, specialists and the authorities. It is now three years since the European Society of Cardiology (ESC), in collaboration with the EASD, published its guidelines on diabetes, pre-diabetes and cardiovascular diseases (10). These guidelines obviously have a different and broader approach to cardiovascular diseases than the aforementioned recommendations, but they are largely consistent with them. However, they go further than the EASD and ADA's guidelines in omitting metformin as the medication of first choice for treating hyperglycaemia in people with established or a high risk of cardiovascular disease. They also include more patients in the latter category.

«We believe that a broad and open debate is now needed on the use of and funding for such medication (semaglutide and tirzepatide)»

Heart failure and chronic kidney disease are common major complications of type 2 diabetes and lead to increased mortality and high costs (13,14). Consequently, the guidelines for these conditions have also been updated (11,15). Perhaps the most important

recommendation in the guidelines for chronic kidney disease is the clear advice to screen for the condition through regular measurements of the urinary albumin/creatinine ratio (UACR) in addition to estimated glomerular filtration rate (eGFR) in all patients with type 2 diabetes. In relation to pharmacological treatment, the Kidney Disease: Improving Global Outcomes (KDIGO) organisation draws attention to new evidence related to the use of SGLT2 inhibitors and non-steroidal mineralocorticoid receptor antagonists in addition to standard treatment with angiotensin-converting enzyme (ACE) inhibitors or angiotensin II receptor antagonists.

Treatment recommendations in Norway

The Norwegian Directorate of Health issued national clinical guidelines for diabetes in 2016 ([16](#)), and these were most recently updated in 2019. Chapters 5, 6 and 8 of the guidelines cover treatment with anti-hyperglycaemic drugs and medications aimed at preventing late sequelae of macrovascular disease as well as kidney disease in people with diabetes. Some of these recommendations deviate significantly from what is set out above. This is a major challenge for doctors who want to treat patients based on the best available evidence base. Explaining treatment options to patients who follow international developments can also be difficult.

Can doctors in Norway not just follow the international recommendations? The main challenge is that approved indications for the various medicines are not always in accordance with updated guidelines and current reimbursement rules.

Cost-benefit analysis

The knowledge underlying the recommendations referred to above is based on large-scale, high-quality, randomised controlled trials, where the outcomes tend to be reported in terms of the relative risk reduction for a primary endpoint. The outcomes obviously vary between trials, but a relative risk reduction of 15–40 % is often seen for a serious cardiac, vascular or renal event.

These are impressive figures, but because serious events are (fortunately) relatively rare during the 2–4-year-long trials, it means that the absolute event rate is likely to be reduced by just 1–3 % and that 20–80 patients must therefore be given the relevant treatment for 2–4 years to prevent just one serious event. Some would consider the cost-benefit impact to be too low, representing too great a burden for the individual patient and an unacceptably high cost for society. We also have to consider that these are normally lifelong treatments, not just something that the patient takes for 2–4 years, and that the events are very serious (stroke, heart attack, cardiovascular death, heart failure and terminal kidney failure). Although there is no strong evidence to support this, it is reasonable to assume that the benefit will increase with the duration of treatment.

What is the current practice in Norway?

The Norwegian Medicines Agency's recommendations regarding approved indications and reimbursement rules are constantly changing, and so is the price of pharmacological agents. It is hoped that indications, reimbursement rules and national treatment recommendations will be brought more in line with the frontier of knowledge that the international clinical guidelines represent.

«It is hoped that indications, reimbursement rules and national treatment recommendations will be brought more in line with the frontier of knowledge that the international clinical guidelines represent»

However, we will have to accept the major discrepancy for the time being, and probably for a long time to come. Some doctors might treat patients without an approved indication, and some patients are already paying for treatment that is not covered by the State. This could lead to a detrimental increase in health inequalities, and we believe it is important that the gap between the updated knowledge base and Norwegian recommendations and actual treatment does not become too wide.

CONCLUSION

We believe doctors in Norway should be able to prescribe an SGLT2 inhibitor with a demonstrated protective effect on the heart and kidneys to patients with type 2 diabetes and known heart, vascular or kidney disease, regardless of their HbA_{1c} level. The same should probably apply to GLP1 receptor agonists for patients with obesity, although the costs here are higher and the cost-benefit impact more uncertain. Perhaps it should also be possible to prescribe these two drug types to patients with a high risk of heart, vascular or kidney disease, but it may be hard to draw a line between moderate and high risk. Doctors should inform patients with type 2 diabetes that significant weight reduction can lead to remission of diabetes, and they should have access to tools that help them achieve this.

REFERENCES

1. McGuire DK, Shih WJ, Cosentino F et al. Association of SGLT2 Inhibitors With Cardiovascular and Kidney Outcomes in Patients With Type 2 Diabetes: A Meta-analysis. *JAMA Cardiol* 2021; 6: 148–58. [PubMed][CrossRef]
2. Sattar N, Lee MMY, Kristensen SL et al. Cardiovascular, mortality, and kidney outcomes with GLP-1 receptor agonists in patients with type 2 diabetes: a systematic review and meta-analysis of randomised trials. *Lancet Diabetes Endocrinol* 2021; 9: 653–62. [PubMed][CrossRef]
3. Palmer SC, Tendal B, Mustafa RA et al. Sodium-glucose cotransporter protein-2 (SGLT-2) inhibitors and glucagon-like peptide-1 (GLP-1) receptor agonists for type 2 diabetes: systematic review and network meta-analysis of randomised controlled trials. *BMJ* 2021; 372: m4573. [PubMed][CrossRef]
4. Lean ME, Leslie WS, Barnes AC et al. Primary care-led weight management for remission of type 2 diabetes (DiRECT): an open-label, cluster-randomised trial. *Lancet* 2018; 391: 541–51. [PubMed][CrossRef]
5. Hofsø D, Fatima F, Borgeraas H et al. Gastric bypass versus sleeve gastrectomy in patients with type 2 diabetes (Oseberg): a single-centre, triple-blind, randomised controlled trial. *Lancet Diabetes Endocrinol* 2019; 7: 912–24. [PubMed][CrossRef]
6. Lingvay I, Sumithran P, Cohen RV et al. Obesity management as a primary treatment goal for type 2 diabetes: time to reframe the conversation. *Lancet* 2022; 399: 394–405. [PubMed][CrossRef]
7. Davies M, Færch L, Jeppesen OK et al. Semaglutide 2.4 mg once a week in adults with overweight or obesity, and type 2 diabetes (STEP 2): a randomised, double-blind, double-dummy, placebo-controlled, phase 3 trial. *Lancet* 2021; 397: 971–84. [PubMed][CrossRef]
8. Rosenstock J, Wysham C, Frías JP et al. Efficacy and safety of a novel dual GIP and GLP-1 receptor agonist tirzepatide in patients with type 2 diabetes (SURPASS-1): a double-blind, randomised, phase 3 trial. *Lancet* 2021; 398: 143–55. [PubMed][CrossRef]
9. Davies MJ, Aroda VR, Collins BS et al. Management of hyperglycaemia in type 2 diabetes, 2022. A consensus report by the American Diabetes Association (ADA) and the European Association for the Study of Diabetes (EASD). *Diabetologia* 2022; 65: 1925–66. [PubMed][CrossRef]
10. Cosentino F, Grant PJ, Aboyans V et al. 2019 ESC Guidelines on diabetes, pre-diabetes, and cardiovascular diseases developed in collaboration with the EASD. *Eur Heart J* 2020; 41: 255–323.

[PubMed][CrossRef]

11. de Boer IH, Khunti K, Sadusky T et al. Diabetes Management in Chronic Kidney Disease: A Consensus Report by the American Diabetes Association (ADA) and Kidney Disease: Improving Global Outcomes (KDIGO). *Diabetes Care* 2022; 45: 3075–90. [PubMed][CrossRef]
 12. National Institute for Healthcare and Care Excellence. Type 2 diabetes in adults: management. <https://www.nice.org.uk/guidance/ng28> Accessed 10.10.2022.
 13. Birkeland KI, Bodegard J, Eriksson JW et al. Heart failure and chronic kidney disease manifestation and mortality risk associations in type 2 diabetes: A large multinational cohort study. *Diabetes Obes Metab* 2020; 22: 1607–18. [PubMed][CrossRef]
 14. Sundström J, Bodegard J, Bollmann A et al. Prevalence, outcomes, and cost of chronic kidney disease in a contemporary population of 2.4 million patients from 11 countries: The CaReMe CKD study. *Lancet Reg Health Eur* 2022; 20: 100438. [PubMed][CrossRef]
 15. McDonagh TA, Metra M, Adamo M et al. 2021 ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure. *Eur Heart J* 2021; 42: 3599–726. [PubMed][CrossRef]
 16. Helsedirektoratet. Nasjonal faglig retningslinje for diabetes. <https://www.helsedirektoratet.no/tema/diabetes> Accessed 10.10.2022
-

Publisert: 20 February 2023. Tidsskr Nor Legeforen. DOI: 10.4045/tidsskr.22.0672

Received 19.10.2022, first revision submitted 16.11.2022, accepted 21.11.2022.

© Tidsskrift for Den norske legeforening 2023. Downloaded from tidsskriftet.no 4 June 2023.