
The term functional disorders should be revitalised

LANGUAGE COLUMN

HELENE HELGELAND

helene.helgeland@gmail.com

Helene Helgeland, PhD, specialist in child and adolescent psychiatry and head of the National Advisory Unit on Complex Psychosomatic Disorders in Children and Adolescents, Oslo University Hospital.

The author has completed the ICMJE form and declares no conflicts of interest.

LINN OKKENHAUG GETZ

Linn Okkenhaug Getz, MD, professor at the Department of Public Health and Nursing, Norwegian University of Science and Technology (NTNU), head of the Research Unit for General Practice in Trondheim, and part of the project management team for the *Sustainability in general practice* collaboration.

The author has completed the ICMJE form and declares no conflicts of interest.

LINN BREEN HERNER

Linn Breen Herner, clinical psychologist at the Department of Child and Adolescent Mental Health in Hospital, Oslo University Hospital, and vice chair of the Committee for Clinical Health Psychology in the Norwegian Psychological Association.

The author has completed the ICMJE form and declares no conflicts of interest.

ERLEND BØEN

Erlend Bøen, PhD, specialist in neurology and psychiatry, senior consultant at the Unit for Psychosomatics and CL Psychiatry, Oslo University Hospital.

The author has completed the ICMJE form and declares the following conflict of interest: he has received lecture fees from Lundbeck.

SVERRE MYREN

Sverre Myren, PhD, specialist in neurology, senior consultant at the Department of Neurology, Ålesund Hospital, researcher at the Department of Neuromedicine and Movement Science, Norwegian University of Science and Technology (NTNU), and associate professor at the Department of Health Sciences, Ålesund, NTNU.

The author has completed the ICMJE form and declares no conflicts of interest.

A new definition of 'functional disorder' refers to changes to the functioning of the nervous system and other organ systems. This understanding has now gained wider acceptance among patients and the medical community, but the continued use of outdated terminology is still causing confusion and misunderstandings.




Illustration: Mimomy/iStock. Edited by the Journal of the Norwegian Medical Association

In medicine, the word *functional* has been used to describe somatic symptoms when standard investigations have failed to detect a pathologic basis in the traditional biomedical sense [\(1–3\)](#). Such symptoms are commonplace, accounting for approximately one-third of consultations in both primary care and the specialist health service [\(4,5\)](#). Examples include headaches, gastrointestinal discomfort, dizziness, fatigue, difficulty concentrating, seizures, paralysis and sensory disturbances. The severity ranges from mild and transient to severe chronic conditions, affecting 1–2 % of the population [\(6,7\)](#).

Terms throughout history

Throughout history, a confusing array of closely related, sometimes overlapping terms have been used to describe such symptoms and conditions. Interestingly, the term *functional* was originally used in the early nineteenth century to specifically describe alterations in the functioning of the nervous system [\(8\)](#). Over the course of the nineteenth century, the meaning of the word evolved into an antonym for organic, as a way of explaining somatic symptoms without a demonstrable physical cause.

At the same time, women with non-specific bodily and emotional symptoms were often diagnosed with *hysteria*, a term rooted in antiquity, where it was believed the symptoms stemmed from a wandering uterus. *Hysterical neurosis* evolved into a psychoanalytic term for bodily symptoms arising from underlying psychological conflicts, meaning they were *psychogenic* in nature. The diagnosis was included in early versions of the ICD (International Classification of Diseases) and DSM (Diagnostic and Statistical Manual of Mental Disorders) until the 1970s.

In the late nineteenth century, Sigmund Freud (1856–1939) and Pierre Janet (1859–1947) introduced the concepts of *conversion* and *dissociation* as defence mechanisms against psychological stress. In conversion, psychological stress was 'converted' into physical symptoms, while dissociation entailed changes in consciousness and memory with a temporary 'disconnect' from reality, thoughts, feelings or sense of identity. These terms were incorporated into international diagnostic classification systems, initially as subcategories of hysterical neurosis. *Somatoform disorders*, which also refer to physical symptoms with psychological origins, was introduced in DSM-III (1980) and ICD-10 (1992, still used in Norway). This category was intended to be more precise and less burdened by the historical and cultural connotations of hysteria.

The term *psychosomatic* has a long tradition and has been used both as an umbrella term to describe how psychological factors and processes affect the body's physiology and to characterise symptoms without a demonstrable physical explanation [\(9\)](#). The term *medically unexplained symptoms* has been in use since the 1990s. It was introduced to reflect the fact that some symptoms may not necessarily have a clear biomedical cause but are still real and can have a major impact on a patient's quality of life [\(10\)](#).

No longer medically unexplained

Research in recent decades has provided deeper insights into causes and mechanisms, showing that they are no longer considered 'medically unexplained'. Cumulative biopsychosocial stress, genetic factors and epigenetic changes impact susceptibility to developing the symptoms. Studies using advanced methods have revealed changes in brain networks and the body's physiology [\(11,12\)](#). There are strong indications that these may be closely related conditions or subcategories with many shared underlying factors [\(5,13,14\)](#).

«It has been proposed that the term functional disorders be used as an umbrella term that encompasses various physical symptoms»

Such symptoms can arise from any of the body's organ systems and are therefore represented in nearly all medical specialties. Accordingly, it has been proposed that the term *functional disorders* be used as an umbrella term that encompasses various physical symptoms (Box 1) (5,15). For example, the international gastroenterology community has, since the 1990s, been developing recognised symptom-based criteria, known as the Rome criteria, for functional gastrointestinal conditions (16).

Box 1 List of diagnostic codes for functional disorders in the Danish ICD-10

DR688A9 Functional disorder NOS
DR688A9A Functional disorder, multiorgan
DR688A9B Functional disorder, single organ
DR688A9B1 Functional disorder, general/fatigue
DR688A9B2 Functional disorder, gastrointestinal
DR688A9B3 Functional disorder, musculoskeletal
DR688A9B4 Functional disorder, cardiopulmonary
DR688A9B5 Functional disorder, neurological
DR688A9B6 Functional disorder, urogenital
DR688A9B9 Other functional disorder, single organ
DR688A9C Functional disorder, single symptom

Today's challenge

Today's challenge is that many outdated terms are still in use – across diagnostic classification systems, among healthcare personnel and within the general population. According to ICD-10, which is used in Norway, several functional symptom profiles are classified under the main categories of somatoform disorders and dissociative (conversion) disorders. However, many clinicians and patients find that terms such as *conversion*, *dissociation* and *somatoform* are not well defined, leading to vague and inconsistent use. In contrast, *functional disorder* is not an established diagnostic category in our classification system but is widely used in clinical practice – primarily in line with the traditional understanding.

Common to all these terms, including *functional* as used in the traditional definition, is that they reflect something medically unexplained and psychogenic. This can give the impression that what the patient is experiencing is not fully understood and may be perceived as an indication that the symptoms are not real and just 'in their head'. Use of these terms also perpetuates an artificial, reductionist dichotomy between body and mind, which is incompatible with modern medical knowledge and understanding.

The euphemistic treadmill

To counteract stigmatisation and this false dichotomy, new terms have been continually introduced throughout history. Many have advocated for discarding terms such as *somatoform*, *psychogenic*, *medically unexplained* and *functional*, while introducing terms like *complex symptom disorders* (a term proposed for use in Norway) (17,18). Others propose terms that specify what the patient's symptoms are *not*, for example, *non-epileptic seizures* and *non-cardiac chest pain* (19). However, whether these represent anything other than superficial changes and euphemisms is debatable. The term *euphemistic treadmill* is used to describe the constant search for new words to replace those that are taboo, only for the new words themselves to eventually be superseded because they too acquire negative connotations.

«Euphemistic treadmill is used to describe the constant search for new words to replace those that are taboo, only for the new words themselves to eventually be superseded because they too acquire negative connotations»

Providing an acceptable explanation for and naming a patient's symptoms are key components of the diagnostic and therapeutic process. However, name changes alone cannot address the fundamental challenges associated with understanding and treating complex functional symptoms; they need to be accompanied by a deeper and more comprehensive shift in how healthcare personnel and society at large perceive these conditions.

Finding terms that are both unambiguous and comprehensive is a challenge in many areas of medicine (5,20). Clinical investigations often conclude with phrases like 'no cause found', or descriptions of the condition as 'complex', or explanations of the patient's symptoms based on what they are *not*. Such practices often become an unfortunate endpoint in the diagnostic process and fail to provide patients with a meaningful explanation for their symptoms. Many patients are left feeling offended, dismissed and disbelieved, leading to misunderstandings and a lack of trust in the patient-doctor relationship (21–23).

A pragmatic solution

The challenges associated with finding terms for medical conditions are well-known and have been the subject of debate for decades. Finding a solution that garners broad consensus has proved to be difficult. However, reputable medical communities in Europe, the United States, Australia and the Nordic countries have now agreed on the term *functional*, but with a new definition: an altered function of the nervous system and other organ systems (5,11,24). This updated definition paves the way for a biopsychosocial understanding of the disease and reflects the natural reversibility of the symptoms.

«Studies and clinical experience indicate that patients perceive the term 'functional' as neutral and inoffensive»

Studies and clinical experience indicate that, contrary to what many doctors believe, patients perceive the term 'functional' as neutral and inoffensive when it is explained to them (22,23). The term is also well-known among medical personnel. Based on this, Denmark introduced its own diagnostic codes for functional disorders in its 2019 version of ICD-10 (25). The diagnostic codes were grouped into a neutral chapter, separate from the chapters for mental disorders and diseases of specific organs (Box 1). This is a pragmatic solution to bringing together many of the partially overlapping terms currently in use. It also aligns with developments in DSM-5 and ICD-11, both of which have introduced functional neurological disorder as a separate diagnosis (11).

The word *functional* should be revitalised

Universal terminology and nomenclature are essential for ensuring clarity, precision and quality in diagnostics, treatment, research and communication within the field of functional disorders. Agreeing on a common international term across medical disciplines is challenging but necessary. We support the work initiated by leading international medical communities. Revitalising the term *functional* is an important step in the right direction.

REFERENCES

1. Barsky AJ, Borus JF. Functional somatic syndromes. *Ann Intern Med* 1999; 130: 910–21. [PubMed][CrossRef]
2. Det Norske Akademis Ordbok. Funksjonell. <https://naob.no/ordbok/funksjonell> Accessed 4.2.2024.
3. Persvold AZ. Funksjonell. <https://snl.no/funksjonell> Accessed 27.11.2024.
4. Myren S, Helgeland H. Legg vekt på positive tegn ved diagnostisering av funksjonell nevrologisk lidelse. *Tidsskr Nor Legeforen* 2024; 144. doi: 10.4045/tidsskr.24.0321. [PubMed][CrossRef]
5. EURONET-SOMA Group. Functional somatic disorders: discussion paper for a new common classification for research and clinical use. *BMC Med* 2020; 18: 34. [PubMed][CrossRef]
6. Creed F, Barsky A, Ann Leiknes K. Epidemiology: prevalence, causes and consequences. Cambridge: Cambridge University Press, 2011: 1–42.
7. Garralda EM, Rask CU. Somatoform and related disorders. *Rutter's Child and Adolescent Psychiatry*. 6. utg. Oxford: Wiley-Blackwell, 2015: 1035–54.
8. Trimble MR. Functional diseases. *Br Med J (Clin Res Ed)* 1982; 285: 1768–70. [PubMed][CrossRef]

9. Malt UF, Malt EA, Blomhoff S et al. Funksjonelle somatiske lidelser—en oversikt. *Tidsskr Nor Lægeforen* 2002; 122: 1379–84. [PubMed]
10. Creed F, Guthrie E, Fink P et al. Is there a better term than "medically unexplained symptoms"? *J Psychosom Res* 2010; 68: 5–8. [PubMed][CrossRef]
11. Aybek S, Perez DL. Diagnosis and management of functional neurological disorder. *BMJ* 2022; 376: o64. [PubMed][CrossRef]
12. Reme SE. Medically unexplained symptoms explained. *Lancet* 2024; 403: 2568–9. [PubMed][CrossRef]
13. Hallett M, Aybek S, Dworetzky BA et al. Functional neurological disorder: new subtypes and shared mechanisms. *Lancet Neurol* 2022; 21: 537–50. [PubMed][CrossRef]
14. Budtz-Lilly A, Schröder A, Rask MT et al. Bodily distress syndrome: A new diagnosis for functional disorders in primary care? *BMC Fam Pract* 2015; 16: 180. [PubMed][CrossRef]
15. Velazquez-Rodriguez Ç, Fehily B. Functional neurological disorder: Historical trends and urgent directions. *J Neurol Res* 2023; 13: 12–32. [CrossRef]
16. Drossman DA. Functional gastrointestinal disorders: History, pathophysiology, clinical features, and Rome IV. *Gastroenterology* 2016; 150: 1262–79.e2. [PubMed][CrossRef]
17. Borchgrevink P, Fredheim O, Fors E et al. Hva er sammensatte lidelser? *Tidsskr Nor Legeforen* 2009; 129: 1368. [CrossRef]
18. Finset A. Klinisk helsepsykologisk kartlegging ved sammensatte symptomtilstander. *Tidsskr Nor Psykol foren* 2024; 61: 915–23.
19. Nakken KO, Villagran A. Fra hysteroepilepsi til ikke-epileptiske anfall. *Tidsskr Nor Legeforen* 2021; 141. doi: 10.4045/tidsskr.20.0832. [CrossRef]
20. Fors EA, Jacobsen HB, Borchgrevink PC et al. Somatiske symptomer uten vesentlig medisinsk forklaring: Bakgrunn, nosologi og kliniske betraktninger med relevans for psykologer. *Tidsskr Nor Psykol foren* 2014; 51: 365–77.
21. Kozłowska K, Sawchuk T, Waugh JL et al. Changing the culture of care for children and adolescents with functional neurological disorder. *Epilepsy Behav Rep* 2021; 16. doi: 10.1016/j.ebr.2021.100486. [PubMed][CrossRef]
22. Stone J, Wojcik W, Durrance D et al. What should we say to patients with symptoms unexplained by disease? The "number needed to offend". *BMJ* 2002; 325: 1449–50. [PubMed][CrossRef]
23. Mcloughlin C, Lee WH, Carson A et al. Iatrogenic harm in functional neurological disorder. *Brain* 2025; 148: 27–38. [PubMed][CrossRef]
24. Kozłowska K, Scher S, Helgeland H. Functional somatic symptoms in children and adolescents: The Stress-System Approach to assessment and treatment. London: Palgrave Macmillan, 2020.

25. Sundhedsstyrelsen. Kodning af funktionelle lidelser - Vejledning til sundhedspersonale. <https://www.sst.dk/da/udgivelser/2020/Vejledning-til-anvendelse-af-koder-for-funktionelle-lidelser> Accessed 6.12.2024

Publisert: 30 January 2025. Tidsskr Nor Legeforen. DOI: 10.4045/tidsskr.24.0573

Received 1.11.2024, first revision submitted 4.12.2024, accepted 6.12.2024.

Copyright: © Tidsskriftet 2026 Downloaded from tidsskriftet.no 13 July 2026.