

Appendix to Ane Sørli Kværner, Helene Harnæs, Dena Helene Alavi, Linnea Bärebring, Hege Berg Henriksen, Marianne Grønlie Guren, Peter Mæhre Lauritzen, Heidi Beate Eggesbø, Gro Wiedswang, Sigbjørn Smeland, Rune Blomhoff. Should calculation of chemotherapy dosage for bowel cancer be based on body composition? Tidsskr Nor Legeforen 2020; 140. doi: 10.4045/tidsskr.19.0769. This appendix is a supplement to the article and has not undergone editorial revision.

Appendix 1: Search strategy

Database: PubMed

Population (((((((("Colorectal Neoplasms"[Mesh:noexp]) OR "Colonic Neoplasms"[Mesh])) OR ((colorectal cancer[Title/Abstract] OR colorectal neoplasm*[Title/Abstract] OR colonic neoplasm*[Title/Abstract] OR colon cancer))))))

AND

Setting ("Antineoplastic Agents"[Mesh] OR adjuvant chemotherapy OR folfox OR folfox protocol OR oxaliplatin OR 5-fluorouracil OR capecitabine))

AND

Exposure ((malnutrition OR nutritional status OR weight loss OR body composition OR sarcopenia OR BMI OR "muscle mass" OR nutritional assessment OR anthropometry OR undernourished OR undernourishment OR wasting))

AND

Outcome (toxicity OR toxic OR toxicities OR adverse reactions OR adverse events OR side effects OR adverse effects OR adverse effect OR side effect OR adverse effect OR dose reduction OR drug-related side effects OR dose modification OR toxicity tests)
